

第01回

- 講演者： **Wolfram Bauer** 氏（学振□□□）
 - 題目□ Hankel operators on the Segal-Bargmann space and Ψ^* -algebras by commutator methods
 - 日時：平成18年4月13日（木）16:00～17:00

For the Segal-Bargmann space $H^2(\mathbb{C}^n, \mu)$ of Gaussian square integrable entire functions on $\mathbb{C}^n$ we consider Hankel operators H_f with f in a symbol space $\mathcal{T}(\mathbb{C}^n)$. We define the Berezin transform for operators on $H^2(\mathbb{C}^n, \mu)$ and in terms of the {itshape mean oscillation} of f we give necessary and sufficient conditions for H_f and $H_{\bar{f}}$ to be bounded, compact or to belong to the {itshape von Neumann Schatten class} $\mathcal{S}_p$ for $1 \leq p < \infty$. We compare some aspects of these results to the case of Bergman spaces over bounded symmetric domains.

There are close relations to the boundedness of the commutator $[P, M_f]$ where P denotes the Toeplitz projection and M_f is the multiplication by f . We describe how to construct spectral invariant Fréchet operator algebras $(\Psi_k^{\Delta})_k$ with prescribed properties in $\mathcal{L}(H^2(\mathbb{C}^n, \mu))$ similar to the Hörmander classes $\Psi_{\rho, \delta}^0$ of zero-order pseudo-differential operators. Following general ideas by B. Gramsch we are using commutator methods and finite systems Δ of closed operators. With Δ in a class of vector fields on $\mathbb{C}^n$ this construction leads to algebras localized in cones $\mathcal{C} \subset \mathbb{C}^n$ and containing the Segal-Bargmann projection.



.lg-outer.lg-pull-caption-up.lg-thumb-open .lg-sub-html {bottom:80px;}

6 images

[<6>]

[PDF bauer.pdf](#)

From:

<https://wiki.ma.noda.tus.ac.jp/rs/> - (旧)理工学部 数学科

Permanent link:

<https://wiki.ma.noda.tus.ac.jp/rs/seminar/2006/001>

Last update: **2017/11/17 17:26**

