# 第13回

- 講演者： **澤田 伸晴** 氏（東京理科大学理工学部)
  - 題目：Polyhedral realization of crystal base $B(\infty)$ for $C^{(1)}_n$
  - 日時：平成21年2月27日(金) 16:00〜17:00

The crystal base has been realized by several methods but it is not so easy to obtain the explicit form. Polyhedral realization of crystal bases is one method for explicity realizing crystal bases, which was introduced by Nakashima and Zelevinsky. We can describe a vector in the crystal base $B(\infty)$ as a lattice point of certain convex polyhedron in an infinite $\mathbb{Z}$-lattice using this method. This method can be applied to the crystal base $B(\lambda)$ of the irreducible integrable highest weight module for symmetrizable Kac-Moody Lie algebras.

The polyhedral realizations of $B(\infty)$ are given when $\mathfrak{g}$ is arbitrary of rank $2$, when $\mathfrak{g}$ is of type $A_n$ and when $\mathfrak{g}$ is of type $A^{(1)}_{n-1}$ and Nakashima gave the polyhedral realizations of the crystal base $B(\infty)$ $(\lambda \in P_{+})$ of the irreducible integrable highest weight module when $\mathfrak{g}$ is the same cases as the above.

Hoshino obtain the polyhedral realizations of $B(\infty)$ and $B(\lambda)$ for $B_n$, $C_n$, $D_n$, $E_6$, $E_7$, $E_8$, and $F_4$. In my talk, we give the polyhedral realization of $B(\infty)$ for $C^{(1)}_n$ (Joint work with Hoshino and Nakada).

[<6>]

From:
https://wiki.ma.noda.tus.ac.jp/rs/ - **(旧)理工学部 数学科**

Permanent link:
**https://wiki.ma.noda.tus.ac.jp/rs/seminar/2008/013**

Last update: **2017/11/17 14:24**